UNIVERSITY OF CALIFORNIA, DAVIS

BERKELEY • DAVIS • IRVINE • LOS ANGELES • MERCED • RIVERSIDE • SAN DIEGO • SAN FRANCISCO

VETERINARY GENETICS LABORATORY SCHOOL OF VETERINARY MEDICINE ONE SHIELDS AVENUE DAVIS, CALIFORNIA 95616-8744



SANTA BARBARA • SANTA CRUZ

TELEPHONE: (530) 752-2211 FAX: (530) 752-3556

HORSE COAT COLOR TEST RESULTS

| MELANIE PETER MOORBURGER ELBDEICH 319 21079 HAMBURG GERMANY | | | | Case: Date Received: Report Date: Report ID: | DT23962 31-Mar-2010 02-Apr-2010 2959-3854-7071-6048 |
|---|--|---------------|----------|--|--|
| | | SMOKEY CADILL | U | | |
| YOB: 10 I | Breed: QH | SEX: S | Alt. ID: | | |
| Dam: SMOKEY CADILLAC Reg: | | | | | |
| RED FACTOR | Both black and red factors detected. Either E or e transmitted to of other modifying genes. | | | offspring. Basic color is b | olack, bay or brown in the absence |
| E/e | | | | - | · |
| AGOUTI | Black pigment distributed in points pattern. Basic color is bay or brown in the absence of other modifying genes. | | | | |
| A/A | | | | | |
| CREAM DILUTION | Heterozygous, dilute, one copy of Cream gene. Typical colors are palomino, buckskin and smoky black in the absence of other modifying genes. | | | | |
| N/Cr | | | | | |
| PEARL DILUTION | Not requested. | | | | |
| SILVER DILUTION | Not requested. | | | | |
| LETHAL WHITE OVERO | Not requested. | | | | |
| SABINO 1 | Not requested. | | | | |
| TOBIANO | Not requested. | | | | |
| CHAMPAGNE | Not requested. | | | | |

Horse Coat Color Results with Explanations

Red Factor

e/e - Only the red factor detected. Basic color is sorrel or chestnut in the absence of other N/N - No evidence of the altered sequence detected. modifying genes.

E/e - Both black and red factors detected. Either E or e transmitted to offspring. Basic color is black, bay or brown in the absence of other modifying genes.

E/E - No red factor detected. Horse cannot have red foals regardless of the color of mate. Basic color is black, bay or brown in the absence of other modifying genes.

Agouti

A/A - Black pigment distributed in points pattern. Basic color is bay or brown in the absence of other modifying genes.

A/a - Black pigment distributed in points pattern. Basic color is bay or brown in the absence of other modifying genes.

a/a - Only recessive allele detected. Black pigment distributed uniformly. Basic color is black in the absence of other modifying genes.

Cream

N/N - No evidence for the Cream dilution altered sequence detected. Basic color is sorrel or chestnut, bay or black in the absence of other modifying genes.

N/Cr - Heterozygous, dilute, one copy of Cream gene. Typical colors are palomino, buckskin and smoky black in the absence of other modifying genes.

Cr/Cr - Double dilute (two copies of Cream gene). Typical colors are cremello, perlino and smoky cream in the absence of other modifying genes.

Pearl

N/N - No evidence of the altered sequence detected.

N/Prl - One copy of the altered sequence detected. If Cream dilution is also present, a pseudo-double Cream dilute phenotype will result.

Prl/Prl - Two copies of the altered sequence detected. On a chestnut base color, a uniform apricot color of body hair, mane and tail will result.

Tobiano

N/N - No evidence of altered sequence detected. Horse is not Tobiano.

N/TO - One copy of altered sequence. Approximately 50% of the offspring will inherit Tobiano.

TO/TO - Two copies of altered sequence. Horse is homozygous for Tobiano. All offspring will inherit Tobiano.

N/Z - One copy of the altered sequence detected. Black-based horses will be chocolate with flaxen or lightened mane and tail. Bay-based horses will have lightened black pigment on lower legs, mane and tail. No effect on chestnut

Z/Z - Two copies of altered sequence detected. Black-based horses will be chocolate with flaxen or lightened mane and tail. Bay-based horses will have lightened black pigment on lower legs, mane and tail. No effect on chestnut

Lethal White Overo

N/N - No evidence for the altered sequence detected.

N/O - One copy of the altered sequence detected. If bred to another N/O horse, there is a 25% chance of producing a lethal white overo foal. The N/O type has been detected in Paints (including breeding stock), Pintos, Thoroughbreds, Miniatures, Quarter Horses and Tennessee Walking Horses.

O/O - Only the altered sequence in the EDNRB gene detected. This result has only been obtained with samples from lethal white overo foals.

Sabino 1

N/N - No evidence of altered sequence detected.

N/SB1 - One copy of the Sabino 1 gene detected. Horse typically may have 2 or more white legs, blaze, spots or roaning in the midsection and jagged margins around white areas.

SB1/SB1 - Two copies of the Sabino 1 gene detected. Complete or nearly complete white phenotype expected.

Champagne

N/N - No evidence of altered sequence detected.

N/Ch - One copy of the altered sequence detected. Chestnut color (red) is diluted to gold, bay to tan with brown points and black to darker tan with brown points.

Ch/Ch - Two copies of the altered sequence detected. All offspring are expected to be Champagne diluted.

Gray

N/N - No copies of the gray gene. Horse will not turn gray.

N/G - One copy of the gray gene. Horse will turn gray and approximately 50% of offspring will be gray.

G/G - Two copies of the gray gene. Horse will turn gray and all offspring will be gray.